

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/435, 16/18, G01N 33/553	A1	(11) International Publication Number: WO 98/01471 (43) International Publication Date: 15 January 1998 (15.01.98)
(21) International Application Number: PCT/SE97/01164 (22) International Filing Date: 27 June 1997 (27.06.97) (30) Priority Data: 9602677-8 5 July 1996 (05.07.96) SE (71) Applicant (for all designated States except US): AB SANGTEC MEDICAL [SE/SE]; P.O. Box 20045, S-161 02 Bromma (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): BRUNDELL, Jan [SE/SE]; Sveavägen 78, S-113 59 Stockholm (SE). NYBERG, Lena [SE/SE]; Tallbacksvägen 30 B, S-756 45 Uppsala (SE). (74) Agents: BERG, S., A. et al.; H. Albiñs Patentbyrå AB, P.O. Box 3137, S-103 62 Stockholm (SE).		(81) Designated States: AU, BR, CA, FI, HU, IL, JP, NO, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: METHODS FOR DETERMINING THE PRESENCE OF BRAIN PROTEIN S-100 (57) Abstract An assay method for determining the presence of the brain protein S-100 in a clinical sample which uses antibodies directed to epitopes in the region from ser1 to asn38 and from thr82 to glu93 of the amino acid sequence of the β subunit of human S100B is provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS FOR DETERMINING THE PRESENCE OF BRAIN PROTEIN S-100

The present invention relates methods for diagnosis and follow-up of patients with cerebral dysfunction as well as melanoma cancer, by determining the presence of the
5 brain protein S-100. The invention also relates to peptides comprising useful antigenic determinants from S-100 as well as monoclonal antibodies binding to these peptides.

As is known, the nervous system contains a number of proteins unique to its various
10 cellular elements. The cellular disruption of nervous tissue and cells of neural origin, by any pathogenic process, trauma or by neurological diseases, results in the release of normal soluble endogenous cytoplasmic proteins into the cerebral extracellular fluid and ultimately to other body fluids including the cerebrospinal fluid (CSF) and blood (serum and plasma). Examples of representative soluble small molecule
15 weight proteins of this type can be found in the S100 protein family. A review of this family can be found in Zimmer et al., Brain Research Bulletin, Vol. 37, pp 417-429, 1995.

Following disruption of cell membranes, these proteins are released into the
20 extracellular fluid in accordance with a time course and in quantities relative to the pathogenesis of the disease process or the extent of the brain tissue damages. The proteins diffuse into the CSF and then the blood or directly into the blood. The above mentioned cell membrane disruption is reflected by the blood plasma or serum levels of one or more of these antigens and markers. These protein antigens have the
25 advantage of being stable and specific, not only for the brain, but for the cellular components in the brain. By following the relative release of the various nervous system protein antigens, it is possible to deduce the kind of destructive process occurring in the course of neurological diseases and/or the extent of possible brain tissue damages. Information of this type permits the diagnosis, evaluation of severity
30 and rate of progression of the above mentioned diseases and damages.

It is previously known to determine the amount of S-100 polypeptides in a clinical sample. US-A-4 654 313 discloses a radioimmunological assay method for S-100

protein. The patent document does neither mention anything about different kinds of S100-polypeptides nor about on which epitopes the assay method is based. The detection limit is declared to be 0.20 ng/ml but concentrations between 1.5 and 2.5 ng/ml is required in order to have less than 10% false positives. This concentration is rather high. Moreover, in some countries it is not permitted to use radioactive methods in clinical assays.

It is also known to determine S-100 polypeptides by using ELISA-related methods. GB-A-2 109 931 discloses a solid-phase immunoanalysis method comprising the use of enzyme-labelled antigens and particles coated with protein A on which antibodies are bound. S-100 proteins are only mentioned in claim 8 and nothing is revealed about the sensitivity of the method.

JP-A-6/109 734 describes a method suitable for analysing S-100 polypeptides, using a first polyclonal antibody fixed to magnetic particles, and a second labelled polyclonal antibody. The method requires two different enzymes, namely horseradish peroxidase and alkaline phosphatase, and it comprises at least ten consecutive steps. The minimum detection limit is stated to be 0.02 ng/ml for cerebrospinal fluid and 0.06 ng/ml for bovine brain.

The complexity of clinical samples is often a serious problem. An assay method may give excellent results with artificial samples in the laboratory but quite a number of unreliable results might be obtained when the method is tested under clinical conditions. When it comes to immunological assays the problems are often caused by an improper selection of antigenic determinants. One antibody in an assay comprising the use of two different antibodies, may be a hindrance to the other antibody when bound to the antigen to be determined. An improper selection of epitope for an antibody involved in the detection process may result in that the detection group is completely or partially embedded in a protein complex and not available for detection. Different proteins present in the sample might interfere. Moreover, a method comprising many consecutive steps may give uncertain results for complex clinical samples, as the interference possibilities increase with the number of steps and added extra components.

There is always a need for improvement of methods for analysing substances of medical interest in clinical samples. An ideal clinical assay method should be quick, accurate and possible to perform with all types of clinical samples without
5 degeneration of the accuracy for certain types of specimen. It should also require a minimum of extra components. This applies to determination of S-100 polypeptides as well as other substances of medical interest.

Summary of the invention

10

Now it has turned out that by using antibodies directed to epitopes in the region from ser1 to asn38 and from thr82 to glu93 of the amino acid sequence of human S-100 β polypeptide, an improved clinical assay method for determining S-100 polypeptides and particularly the β subunit or isoform thereof is obtained. Hence, the main object
15 of the present invention is an assay method using monoclonal antibodies directed to these epitopes. Another object of the present invention relates to short peptides having sequences corresponding to parts of the amino acid sequence of the human S-100 β polypeptide from ser1 to asn38 and from thr82 to glu93. Yet another object of the present invention relates to analytical kits for carrying out the assay methods.

20

Detailed description of the invention

As already mentioned above it is often very difficult to outline methods for analysing clinical samples. It is necessary that the method has a high sensitivity and
25 gives accurate results. It is also very important that known and unknown constituents of the sample other than the analyte do not influence the results. The present invention relates to an immunological assay method for determining the presence and/or content of human S-100 polypeptide based upon a selection of suitable S-100 epitopes and corresponding antibodies which fulfil the above mentioned
30 requirements.

It has turned out that the selected epitope combinations provides tests and test kits where:

1. a high sensitivity is achieved;
- 5 2. the antibodies of the kit binds equally strong to the internal standard as to the analyte in the clinical sample;
3. the epitopes are chosen in such a way that the different antibodies do not interfere with each other when they bind to the analyte, i.e. that the epitopes are situated sufficiently distant from each other.

10

The epitopes of the present invention are all comprised in the human S-100 β polypeptide. Epitopes present within the amino acid sequences:

SELEKAMVALIDVFHQYSGREGDKHKLKKSELKELINN (SEQ. ID. NO. 2)
and

15 TACHEFFEHE (SEQ.ID.NO. 3)

are preferred. Particularly preferred are epitopes comprised within the peptide AMVALIDVFHQYSGREGDKHKLKKSELKELINN (SEQ. ID. NO. 4) and especially within the peptides REGDKHKLKKSELKEL (SEQ. ID. NO. 5) and EFFEHE (SEQ. ID. NO. 6).

20

The disclosed epitopes are, among all, used to construct peptides for inducing the formation of suitable antibodies on which the claimed assay method is based. These peptides mostly consist of up to 38 amino acids. The whole amino acid sequence of a peptide according to the present invention is derived from human S-100 β . These
25 peptides may comprise variants wherein the original amino acid sequence is modified or altered by insertion, addition, substitution, inversion or deletion which preferably show at least 90% homology with the sequence of SEQ. ID. NO. 2 and SEQ.ID.NO.3 and retain essentially the same immunological properties. The peptides may also comprise multiples of certain epitopes, and in this case their
30 sequence length may exceed 38 amino acids.

By the expression "sub-fragment" is meant a polypeptide sequence having a length of at least 6 amino acids.

The epitopes can also be used to construct fusion peptides comprising at least two distinct epitopes which, among all, can be used as internal standard in immunoassays.

Abbreviations

10 The following abbreviations are used:

	S100	-S100 β
	RT	-Room Temperature
	BSA	-Bovine Serum Albumin
15	Mab(s)	-Monoclonal antibody(ies)
	kD	-kiloDalton
	ECL	-Enhanced Chemiluminescent Assay
	CBB	-Commassie Brilliant Blue
	LIA	-Luminometric Immuno Assay
20	IRMA	-Immuno Radio Metric Assay
	ELISA	-Enzyme Linked ImmunoSorbent Assay
	SDS-PAGE	-SodiumDodecylSulfate - PolyAcrylamideGelElectrophoresis
	PBS	-Phosphate Buffered Saline
	RLU	-Relative Light Units
25	NHS	-N-HydroxySuccinimide
	EDC	-N-ethyl-N'-(dimethylaminopropyl)-carbodiimide
	RAMFc	-RabbitAnti-MouseFc antibody
	EDTA	-EtylenDiamineTetraAcetic acid
	NaCl	-Sodium Chloride
30	NaN ₃	-Sodium azide
	iv.	-intravenously
	aa	-amino acid

	ng	-nanogram
	ml	-millilitre
	mg	-milligram
	HRP	-HorseRadish Peroxidase
5	h	-hour(s)
	min	-minute(s)
	sec	-second(s)

Experimental details common to all test procedures

10

The peptides were prepared by the methods disclosed in Merrifield (1963), J. Am. Chem. Soc., vol. 85, p 2149; Gutte et al.(1971), J. Biol Chem vol. 246, p. 1922; and Carpino et al. (1970), J Am Chem Soc vol. 92, p. 5748.

- 15 The monoclonal antibodies were prepared by the method according to Köhler et al.(1975), Nature vol. 256, p. 495; and Harlow et al.(1988), Antibodies. A Laboratory Manual. Cold Spring Harbor, p. 139.

Antigen and Standard preparations

20

Procedure for preparation and purification of S100 antigen prior to immunisation of Balb/c mice was according to Moore (Biochim. Biophys. Res. Comm. 1965, 19: 739 - 744) with a slight modification according to Haglid & Stavrou (J. Neurochem. 1973, 20:1523-1532). Briefly, bovine brain was homogenised in Tris buffer, pH 7.2.

- 25 The homogenate was centrifuged at 10.000 r.p.m. and the clear supernatant was used for further purification by ammonium sulphate precipitation. The fraction still soluble after saturation by ammonium sulphate was dialysed and purified by separation on a Sephadex G150 Sepharose (Pharmacia Biotech AB, Uppsala Sweden)chromatographic column followed by separation on a DEAE-sephadex
30 (ionic exchange) column (Pharmacia Biotech AB, Uppsala Sweden). The fraction eluted by 0.3 - 0.4 M NaCl was collected, desalted, lyophilised and used for further experiments.

Hybridoma construction.

Balb/c mice were immunised with purified S100 β intraperitoneally in Freund's complete adjuvant and were given booster iv. injection 6 weeks later during 3 consecutive days. The spleen was removed on the fourth day after last injection and prepared for fusion. The myeloma cell line Sp2/0-Ag14 was used for fusion of Balb/c spleen cells.

Antibody purification and subclass determination.

10

Monoclonal antibodies were identified, extracted and purified from hybridoma supernatant according to Harlow & Lane Eds. in ANTIBODIES A LABORATORY MANUAL, Cold Spring Harbour Laboratory Press, New York 298-299 & 311.

Briefly, positive hybridoma clones carrying supernatant with specific antibodies were identified using ELISA with microtitreplate wells coated with S100 β . Immunoglobulins were precipitated using saturated ammoniumsulphate and dialysed against 1.5 M Glycine, 3 M NaCl, pH 8.9. Dialysed material were affinity chromatography purified on an protein-A Sepharose (Pharmacia Biotech AB, Uppsala Sweden) column. Fractions were neutralised by addition of small volumes of 1M Tris pH 8.0.

20

Epitope mapping

S100 β (monomer) epitopes for respective antibody was investigated by use of a synthetic peptide library. Peptides were linked to nitro-cellulose filter membrane via an amide link, according to the manufacturer (Research Genetics, USA) and covers all ninety-one aa in the protein. In total the library consisted of thirty-one, all except one being ten aa-residues long synthetic peptides. Each peptide was consecutively shifted three aa towards the -COOH terminal end of the protein. Positive antibody-binding was indicated by the use of a second anti-mouse antibody conjugated with HRP and detected using an ECL assay (Amersham, UK).

30

Results:

Two binding sequences were found

Epitope 1

AMVALIDVFHQYSGREGDKHKLKKSELKELINN (residues 6-38)(SEQ. ID.
5 NO.4)

and

Epitope 2

10 EFFEHE (residues No 86-91) (SEQ. ID. NO. 6)

Antibody reactivity

15 Purified antibodies reacting with the epitopes were checked for reactivity and
affinity using the BIAcore™ system (Pharmacia Biosensor AB, Uppsala Sweden).
Briefly, in order to test the specificity of the antibodies, the RAMFc was
immobilised onto the sensor chip CM5 NHS-ester activated surface, according to
standard procedure, to provide approximately 600 RLU. Then each Mab was bound
20 to the RAMFc surface to approximately 300RLU, followed by the S100αα and the
S100 standard (consisting of 50% S100αβ and 50% S100ββ) in separate
experiments. All reactions were carried out in continuous flow of the phosphate
buffer. The kinetics between antibodies and antigen was done similarly. S100
antigen was added to the chips at 200-450nM for reactivity measurements of the
25 antibody intended for the solid phase and at 1000-1500nM for measurements of the
antibody intended for tracers. Kinetics was determined using the BIAcore™ Kinetic
evaluation 2.1. software (Pharmacia Biosensor AB, Uppsala Sweden). It can be
concluded from the reactivity profile that the antibodies reactive with the epitopes
are specific for the β-containing forms of S100 and not the α-containing form.

Example 1Development of an immunoluminometric procedure

- 5 Tracer antibody was conjugated with luminol. Briefly, ABEI (Sigma, St Louis, Ms) was linked with a diactivated ester (Etylenglykolbis-succimidyl succinat, EGS). The ABEI-EGS-conjugate was next mixed with monoclonal antiS100-antibody in an approximately 50:5 molar ratio in 100µl of PBS pH 7.4, containing 15% acetonitrile and incubated 1 h at room temperature. The ABEI-conjugated antibody was purified
10 on a Sephacryl®S 300 HR (Pharmacia Biotech AB, Uppsala Sweden) gelfiltration column, and appropriate fractions were pooled and diluted in phosphate-buffer.

Preparation of antibody coated tubes for LIA.

- 15 Polystyrene tubes (Greiner, Germany) were incubated overnight at room temperature with 3µg of S100-antibodies in 300 µl of PBS pH 7.5. The tubes were washed with 0.1% Tween20® in PBS. Next, tubes were blocked with a solution containing 0.9%BSA and 4% Saccarose and incubated for 24h. The solution was aspirated and the tubes allowed to dry.

20

LIA test procedure.

- The test was conducted in a two step procedure by incubating 100µl of patient body fluid in antibody coated tubes, or S100 standard with 100 µl of diluent (PBS +
25 5%BSA) and incubated at room temperature. After washing 200 µl of the luminol-labelled antibody was added and a further 2 h of incubation was performed before measurement. After another washing the luminescence was developed using the LIA-mat starter service kit (Byk-Sangtec, Diezenbach Germany) and immediately measured as integrals over a period of 5 sec in luminometer (Berthold, Germany). In
30 order to convert the obtained light signal into concentration of S100 measurements on patient samples were compared with measurements on solutions with known

concentrations of S100 (standards). The detection limit (zero standard +3 standard deviations) was approximately 0.01 µg/l.

Preparation of a Standard curve

5

S100B protein was obtained from Medisera, Lund, Sweden, and diluted in PBS + 5% BSA. Dilutions contained: 0.10, 0.40, 2.00, 8.00 and 20.00 µg/l, of an S100 preparation consisting of 50 % of the ββ form and 50 % of the αβ form. PBS + 5 % BSA was used as standard 0. Three measurements were carried out for each dilution.

10 The measured results as well as statistical calculations are presented in table 1 below:

Table 1					
	Concentration	Counts	Average	Calculated conc.	Average
15	Standard 0	1996		0 µg/l	
		2024		0 µg/l	
		2053		0.0019 µg/l	
			2024		0 µg/l
20	0.10 µg/l	3142		0.135 µg/l	
		2760		0.0647 µg/l	
		2988		0.105 µg/l	
			2963		0.10 µg/l
25	0.40 µg/l	5494		0.394 µg/l	
		5620		0.405 µg/l	
		5579		0.401 µg/l	
			5564		0.40 µg/l
30	2.00 µg/l	21430		2.049 µg/l	
		21028		1.988 µg/l	
		20869		1.966 µg/l	
			21109		2.00 µg/l

Concentration	Counts	Average	Calculated conc.	Average
5	8.00 µg/l	68389	7.823 µg/l	
		67013	7.677 µg/l	
		74791	8.494 µg/l	
		70064		8.00 µg/l
10	20.00 µg/l	175560	22.12 µg/l	
		155141	18.54 µg/l	
		161052	19.51 µg/l	
		163918		20.00 µg/l

The lower detection limit was defined as three standard 0 determinations plus 3X the standard deviation value. For this measurement, it was calculated to be 0.006 µg/l.

15

Clinical determinations of S100 in serum

The S100 concentration was determined in serum from patients receiving heart bypass surgery and being connected to a heart-lung machine. The results are presented in table 2 below:

Table 2				
Patient	Counts	Average	Concentration	Average
1	94698		10.61 µg/l	
	98104		10.99 µg/l	
		96401		10.80 µg/l
2	1716		Not detected	
	1478		Not detected	
		1597		Not detected
3	3762		0.23 µg/l	
	3799		0.23 µg/l	
		3780		0.23 µg/l

30

		12		
Patient	Counts	Average	Concentration	Average
4	13158		1.04 µg/l	
	14183		1.15 µg/l	
		13670		1.10 µg/l
5	8788		0.66 µg/l	
	8580		0.64 µg/l	
		8684		0.65 µg/l
6	10301		0.78 µg/l	
	10100		0.77 µg/l	
10	10200			0.77 µg/l

Example 2

Development of an ELISA test procedure.

15

As tracer antibody was used monoclonal antiS100 antibody conjugated with β -galactosidase according to Harlow & Lane Eds. in ANTIBODIES A LABORATORY MANUAL. Cold Spring Harbour Laboratory Press, New York page 351.

20 Preparation of antibody coated microtiter wells for ELISA.

Microtiterplatewells (Corning, Denmark) were incubated overnight at +4°C with 2.5µg of microtiter wells were finally washed three times with 0.05% Tween20® and air dried before use.

25

ELISA test procedure.

The ELISA was conducted in a multiple step incubation procedure.

100 µl of 1:1 diluted patient sample or 100µl of S100 standard (0 - 20 µg/ml) was added to the wells.

30

The plate was incubated for 1.5h at RT under shaking.

The plates were washed three times with 300µl 0.05% Tween20® in PBS.

100 µl of alkaline phosphatase conjugated tracer antibody was added and a further 1.5h of incubation on a shaker was performed.

The wells were then washed three times with 0.05% Tween20® in PBS.

- 5 100µl of a 5% o-nitro-phenyl-β-galactoside substrate solution was added and the plates were incubated with substrate for another forty-five minutes and colour is developed.

The colour development was stopped by the addition of 100µl 0.66M Na₂CO₃.

- 10 Each well of the plate was read at 405nm in a standard microtiterplate reader. In order to convert the obtained colour signal into concentration of S100, measurements on patient samples were compared with measurements on solutions with known concentrations of S100 (standards). The detection limit (zero standard +3 standard deviations) was approximately 0.2 µg/l.

- 15 Result:

Standard (µg/l)	0	0.5	1.5	5	15
A 405	0.088	0.147	0.244	0.675	1.196

- 20

Example 3

Development of an immunoradiometric (IRMA) test procedure

- 25 IRMA tracer antibody conjugation

A monoclonal antiS100 antibody was conjugated with Iodine using the Chloramine T method according to Greenwood et al. (Biochem. J. 1963, 89:114-123). The specific activity was determined to be 520±80 MBq/mg

- 30

Preparation of antibody coated to polystyrene beads

Monoclonal anti S100 antibodies were coupled to polystyrene beads by the Glutaraldehyde coupling method according to Harlow & Lane Eds. in ANTIBODIES, A LABORATORY MANUAL, Cold Spring Harbour Laboratory Press, New York, 533 & 536-537. Final blocking was by 1% BSA, 0.1% NaN₃ in PBS pH7.5.

IRMA test procedure.

100µl of patient sample or standard was added to polystyrene tubes together with 100µl PBS diluent. One polystyrene coated bead was added to each tube followed by incubation for 1 h at RT on a shaker. Next the beads were washed once with 2ml of demineralised water and 200µl of I-125 labelled tracer antibody was added and the tubes were incubated a further 2h on a shaker. After washing the radioactive signal on the bead was measured in a standard γ-counter. In order to convert the obtained radioactive signal into concentration of S100, measurements on patient samples were compared with measurements on solutions with known concentrations of S100 (standards). The detection limit (zero standard +3 standard deviations) was approximately 0.1 µg/l.

Example 4

Use of IRMA test procedure for assay of S100 in serum from melanoma patients

The S100 based test procedure was applied on clinical questions relating to melanoma. Blood samples from patients with melanoma of various stages of cancer progression were collected in serum collecting tubes. Samples were then frozen and treated according to the test procedure described above in Example 3.

Results:

Relationship to staging.

Clinical Stage I vs Clinical Stage II. In a study of 577 patients the geometric mean for Stage I was found to be 0.12 µg/l and for Stage II the geometric mean was found to be 0.33 µg/l.

p-value < 0.001.

5

Example 5

Use of IRMA test procedure for assay of S100 in serum from melanoma patients

10 The S100 based test procedure was applied on clinical questions relating to melanoma. Blood samples from patients with melanoma of various stages of cancer progression were collected in serum collecting tubes. Samples were then frozen and treated according to the test procedure described above in Example 3.

15 Results:

Relationship to survival

Clinical Stage I vs Clinical Stage II and III. In a study with respect to survival performed on 643 patients the relative hazard and 95% confidence interval was
20 calculated. The relative hazard was found to be 12.3 and the confidence interval 5.6-27.2 with a p-value of <0.001

Example 6

25 Use of the S100 LIA-method for evaluation of the influence of extra corporal circulation equipment on the brain

The S100 based test procedure in Example 1 was applied on monitoring cerebral injury following extra corporeal circulation (ECC). Blood samples from patients
30 undergoing extra corporeal circulation were collected in serum tubes and treated according " Test procedure". Results

	Before start of ECC	End of ECC	1 day after sur- gery	2 days after sur- gery
S100 levels $\mu\text{g/l}$	0	1,67	0,21	0,13

In this group of patients the level of S100 in serum was elevated for at least 2 days after surgery.

Uncomplicated cases should return to normal levels within the first 24 hours (Ref P.

5 Johnson et al. J. Cardiothor. Vasc. Anaesthesia, 9:6 (1995) 694-99).

Example 7

10 Use of LIA test procedure for assay of S100 in serum from melanoma patients

The S100 based test procedure was applied on clinical questions relating to melanoma. Blood samples from patients with melanoma of various stages of cancer progression and blood donors were collected in serum collecting tubes. Samples
15 were frozen and treated according to the test procedure described above in Example 1.

Result: Of 136 patients with various stages of melanoma 25 had a level of S100 below 0.08 and of 100 blood donors tested on the same occasion 7 had a level equal
20 to or above 0.08 $\mu\text{g/l}$.

Example 8

The reliability of both the test and the S100 β polypeptide marker per se when
25 diagnosing melanoma were investigated. On 252 patients with melanoma. serum was drawn before treatment was started and determination of the level of S100 β polypeptide was performed by the assay method disclosed in example 1. When a cut-off value of 0.16 $\mu\text{g/l}$ was used, the medium survival time of patients having a S100 β concentration above the cut-off value was 7 months, whereas the medium

survival time was more than 120 months for patients having a S100 β concentration below the cut-off value.

5 In a patient diagnosed with malignant melanoma, considered to show no evidence of disease and monitored by the immunoradiometric assay method as disclosed in example 3, elevated levels of S100 β were recorded 2 months prior to the appearance of skin metastases and 6 months before metastases in organs were found.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: AB Sangtec Medical
- (B) STREET: P.O. Box 20045
- (C) CITY: Bromma
- (E) COUNTRY: Sweden
- (F) POSTAL CODE (ZIP): 161 02
- (G) TELEPHONE: +46 8 635 12 00
- (H) TELEFAX: +46 8 29 21 81

(ii) TITLE OF INVENTION: Methods for determining brain antigens

(iii) NUMBER OF SEQUENCES: 8

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 91 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Ser Glu Leu Glu Lys Ala Val Val Ala Leu Ile Asp Val Phe His Gln
1 5 10 15

Tyr Ser Gly Arg Glu Gly Asp Lys His Lys Leu Lys Lys Ser Glu Leu
20 25 30

Lys Glu Leu Ile Asn Asn Glu Leu Ser His Phe Leu Glu Glu Ile Lys
 35 40 45
 Glu Gln Glu Val Val Asp Lys Val Asn Glu Thr Leu Asp Ser Asp Gly
 50 55 60
 Asp Gly Glu Cys Asp Phe Gln Glu Phe Met Ala Phe Val Ala Met Ile
 65 70 75 80
 Thr Thr Ala Cys His Glu Phe Phe Glu His Glu
 85 90

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Ser Glu Leu Glu Lys Ala Met Val Ala Leu Ile Asp Val Phe His Gln
 1 5 10 15
 Tyr Ser Gly Arg Glu Gly Asp Lys His Lys Leu Lys Lys Ser Glu Leu
 20 25 30
 Lys Glu Leu Ile Asn Asn
 35

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Thr	Ala	Cys	His	Glu	Phe	Phe	Glu	His	Glu
1			5					10	

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Ala	Met	Val	Ala	Leu	Ile	Asp	Val	Phe	His	Gln	Tyr	Ser	Gly	Arg	Glu
1			5					10					15		

Gly	Asp	Lys	His	Lys	Leu	Lys	Lys	Ser	Glu	Leu	Lys	Glu	Leu	Ile	Asn
		20					25						30		

Asn

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Arg	Glu	Gly	Asp	Lys	His	Lys	Leu	Lys	Lys	Ser	Glu	Leu	Lys	Glu	Leu
1				5				10					15		

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Glu	Phe	Phe	Glu	His	Glu
1				5	

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Asp	Lys	His	Lys	Leu	Lys	Lys	Ser	Glu	Leu
1				5					10

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Lys	Leu	Lys	Lys	Ser	Glu	Leu	Lys	Glu	Leu
1				5					10

Claims

1. A peptide consisting of at least one sub-fragment of the human S-100 β polypeptide comprising from 6 to 38 amino acids, where said sub-fragments show at
5 least 90% homology with the sequence

SELEKAMVALIDVFHQYSGREGDKHKLKKSELKELINN (SEQ. ID. NO. 2)

and/or the amino acid sequence

10

TACHEFFEHE (SEQ. ID. NO. 3)

and retain essentially the same immunological properties.

- 15 2. A peptide according to claim 1 **characterized** in that the sub-fragments are derived from the amino acid sequence:

SELEKAMVALIDVFHQYSGREGDKHKLKKSELKELINN (SEQ. ID. NO. 2).

- 20 3. A peptide according to claim 2, which is

REGDKHKLKK (SEQ. ID. NO. 5);

DKHKLKKSEL (SEQ. ID. NO. 7); or

KLKKSELKEL (SEQ. ID. NO. 8).

25

4. A peptide according to claim 1, **characterized** in that the sub-fragments are derived from the amino acid sequence:

TACHEFFEHE (SEQ. ID. NO. 3).

30

5. A peptide according to claim 4, which is

EFFEHE (SEQ. ID. NO. 6).

6. A peptide according to claim 1, **characterized** in that it consists of at least one sub-fragment derived from the sequence according to SEQ. ID. NO. 2 and at least
5 one sub-fragment derived from the sequence according to SEQ. ID. NO. 3.

7. A monoclonal antibody or a fragment of such an antibody specifically binding a peptide according to anyone of the preceeding claims.

10 8. A monoclonal antibody or an antibody fragment according to claim 7, specifically binding a peptide according to claim 2.

9. A monoclonal antibody or an antibody fragment according to claim 7, specifically binding a peptide according to claim 4.

15 10. Use of a monoclonal antibody or an antibody fragment according to anyone of claims 7-9 in immunological assay methods.

20 11. Use of a peptide according to anyone of claims 1-6 for eliciting antibodies.

12. Use of a peptide according to anyone of claims 1 - 6 in immunological assay methods.

25 13. A method of determining the presence of human S-100 β polypeptide in a sample comprising the steps of:

letting the sample to be analyzed immunologically react with a first monoclonal antibody according to claim 8, said first antibody being coupled to a carrier;

30 letting the sample immunologically react with a second monoclonal antibody according to claim 9, said second monoclonal antibody being provided with detection means;

Washing; and

detecting the amount of S-100 β polypeptide in the sample.

14. A method according to claim 13 where the detection means is a group having the
5 ability of emitting luminescence.
15. A method according to claim 14, where the carrier is a magnetic particle.
16. A kit for determining the presence of human S-100 β polypeptide in a sample,
10 comprising a peptide according to anyone of claims 1 - 6 and/or an antibody
according to anyone of claims 7 - 9.
17. A kit according to claim 16 comprising a first monoclonal antibody according to
claim 8 and a second monoclonal antibody according to claim 9, said first
15 monoclonal antibody being coupled to a carrier and said second monoclonal
antibody being provided with a detection means.
18. A kit according to claim 17, wherein said carrier is a magnetic particle and said
detection means is a group having the ability of emitting luminescence, such as
20 luminol.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/01164

A. CLASSIFICATION OF SUBJECT MATTER		
IPC6: C07K 14/435, C07K 16/18, G01N 33/553 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
IPC6: C07K, A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
WPI, PAJ, CA, MEDLINE, BIOSIS, DBA, PCI GENBANK/EMBL/SWISSPROT/DBJ		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	The Journal of Biological Chemistry, Volume 263, No 16, June 1988, Linda J. Van Eldik et al, "Synthesis and Expression of a Gene Coding for the Calcium-modulated Protein S100Beta and Designed for Cassette-based, Site-directed Mutagenesis" page 7830 - page 7837 --	1-6
A	Proc. Natl. Acad. Sci., Volume 81, October 1984, Linda J. Van Eldik et al, "Production and characterization of monoclonal antibodies with specificity for the S100Beta polypeptide of brain S100 fractions" page 6034 - page 6038	1-6
X	--	7-12, 16-18
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
29 October 1997		30.10.1997
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86		Authorized officer Patrick Andersson Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/01164

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Protein Science, Volume 4, 1995, Craig Donaldson et al, "Human S100b protein: Formation of a tetramer from synthetic calcium-binding site peptides" page 765 - page 772 --	1-3
X	Dialog Information Services, File 351, (World Patent Index), Dialog accession no. 009890614, WPI accession no. 94-170530/21, SRL KK: "Highly sensitive antigen determin.- comprises solidifying antibody on analyte antigen, blocking solid phase, reacting with sample, reacting with peroxidase", JP,A,6109734, 940422, 9421 (Basic) -----	13-15